29. Congenital heart disease/Kawasaki's disease

| Abstract No. | First Name | Last Name | Abstract Title | Session ID | Language |
|--------------|------------|-----------|---|------------|----------|
| 10075 | Hideo | Ohuchi | Plasma Volume Status and the Unique Associations with Fontan Pathophysiology | 29-1 | English |
| 10394 | Tomoko | Ishizu | The Current Clinical Burden of Unplanned Hospitalization of Adult Congenital Heart Disease in Japan | FRS-3 | 日本語 |
| 10721 | Ichiro | Sakamoto | Leaflet Anomaly is Responsible for Tricuspid Regurgitation in Congenitally Corrected Transposition of the Great Arteries and Systemic Right Ventricle | 29-1 | English |
| 10831 | Shuhei | Toba | Automated phenotyping of congenital heart disease and a tool for the interactive reporting of congenital heart disease outcomes | 29-3 | 日本語 |
| 11039 | Ayako | Ishikita | Extracellular volume measured by computed tomography was correlated with right ventricular fibrosis in patients with adult congenital heart disease | 29-1 | English |
| 11064 | Yumi | Shiina | Secondary sarcopenia assessed by computed tomography can predict hospitalization for heart failure in adults with Fontan circulation | 29-2 | English |
| 11068 | Yumi | Shiina | Aortic Vorticity, Helicity and Aortopathy in Adult Patients with Tetralogy of Fallot: preliminary study using four-dimensional flow magnetic resonance images | 29-1 | English |
| 11074 | Yumi | Shiina | Non-physiological aortic flow and aortopathy in adult patients with transposition of the great arteries after LeCompte procedure | 29-1 | English |
| 11199 | Yuji | Tominaga | Impact of diastolic dysfunction on the outcome of pulmonary valve replacement for pulmonary valve regurgitation | 29-2 | English |
| 11260 | Michiyo | Yamano | The Remodeling Process After Transcatheter Atrial Septal Defect Closure: Comparison Between Patients with Atrial Fibrillation and Sinus Rhythm | 29-2 | English |
| 11364 | Tomoko | Ishizu | Heart Failure in Adult Congenital Heart Disease in Japan; A Report from JROAD - DPC Study | 29-3 | 日本語 |
| 11394 | Makiko | Suzuki | Preoperative Right Ventricular Function and Its Coupling with Pulmonary Vasculature are Determinants of Residual Symptoms after Atrial Septal Defect Closure | 29-2 | English |

29. Congenital heart disease/Kawasaki's disease

| Abstract No. | First Name | Last Name | Abstract Title | Session ID | Language |
|--------------|------------|-----------|---|------------|----------|
| 11718 | Keita | Masuda | Effectiveness and Safety of Direct Oral Anticoagulants in Adult Congenital Heart Disease Patients | 29-3 | 日本語 |
| 11738 | Naoto | Kawamatsu | A high VE/VCO ₂ slope in Adult Patients with repaired Tetralogy of Fallot might show hemodynamic, not ventilatory inefficiency. | 29-3 | 日本語 |
| 11836 | Yasuhiro | Tanabe | Endothelialization of the Closure Device for the Atrial Septal Defect (ASD) 6-months after Implantation Evaluated by Angioscopy. | 29-2 | English |
| 11943 | Norihisa | Toh | Renal Congestion Assessed by Doppler Ultrasonography in Adults with Complex Congenital Heart Disease after Biventricular Repair | 29-1 | English |
| 12141 | Teiji | Akagi | Current Condition of Transcatheter Patent Foramen Ovale Closure in Japan: CVIT ASD/PFO Commitee Report | 29-2 | English |
| 12179 | Akio | Inage | Native hepatic T1 may be a potential marker for liver fibrosis in Fontan patients | 29-1 | English |
| 12213 | Eiji | Yamashita | Prevalence, Clinical Features and Outcome of Interatrial Left to Right Shunt Detected by Transesophageal Echocardiography Prior to Atrial Fibrillation Ablation | 29-3 | 日本語 |
| 12304 | Shunsuke | Tatebe | Current Clinical Characteristics and Outcomes of Patients with Adult Congenital Heart Disease Associated with Pulmonary Hypertension | 29-3 | 日本語 |